




Open Archive Toulouse Archive Ouverte (OATAO)

OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible

This is an author's version published in: <http://oatao.univ-toulouse.fr/22752>

Official URL: <https://doi.org/10.1016/j.nbt.2014.05.1967>

To cite this version:

Rahbany, Gihane and Salameh, Dominique and Brandam, Cédric  and Lteif, Roger *Analysis of the effectiveness of a cereal milling by product monocomponent medium for the low cost production of Bacillus thuringiensis*. (2014) In: 16th European Congress on Biotechnology, 13 July 2014 -16 July 2014 (Edinburgh, United Kingdom)

Any correspondence concerning this service should be sent to the repository administrator: tech-oatao@listes-diff.inp-toulouse.fr

Analysis of the effectiveness of a cereal milling by product monocomponent medium for the low cost production of *Bacillus thuringiensis*

Gihane Rahbany^{1,*}, Dominique Salameh¹, Cedric Brandam², Roger Lteif¹

¹ Université Saint-Joseph Liban

² Université de Toulouse; INPT, UPS; Laboratoire de Génie Chimique France

Bacillus thuringiensis is a facultative anaerobe, gram positive, spore forming bacterium. The biotechnological importance of this bacterium resides in its ability to produce, during sporulation, crystal proteins known as δ -endotoxins which express specific insecticidal activity. At industrial scale, the culture media represents an important part of *B. thuringiensis* based biopesticides production cost. According to the literature, different agro-industrial residues and byproducts were used as sources of proteins in order to reduce the cost of *B. thuringiensis* culture medium, but carbohydrates (glucose, starch or molasses) and/or mineral sources were added.

In this work, a cereal milling by-product (CMB) as a monocomponent medium was investigated and compared to synthetic mediums in terms of δ -endotoxin yield and productivity in submerged fermentation of different strains of *B. thuringiensis*. The CMB was shown efficient to be used as a complete substrate (source of proteins, carbohydrates and minerals) for *B. thuringiensis* production. The optimal CMB ratio in the culture medium was found to be 6% in shake flasks experiments. The consumption of the CMB sugars by the bacteria was analyzed. Production of the bio-insecticide in lab-bioreactor in controlled conditions was equally performed to give basic elements for extrapolation in industrial conditions.

<http://dx.doi.org/10.1016/j.nbt.2014.05.1967>